



EJVES Extra Abstracts[☆]

Failure of Sweat Gland Curettage to Relieve Axillary Hyperhidrosis: A Salutary Lesson

S.J. Chapman, M.J. Gough
*The Leeds Vascular Institute, The General Infirmary at Leeds,
Great George Street, Leeds LS1 3EX, UK*

Introduction: Several treatment options are available for primary hyperhidrosis. Selection for individual patients is influenced by symptom severity, success rates and the relative risk of compensatory hyperhidrosis.

Case report: A 24-year-old female presented with a 10-year history of palmar, pedal and axillary hyperhidrosis. Following recurrent relapse after botulinum toxin (BOTOX[®]) injections for axillary hyperhidrosis bilateral axillary sweat gland curettage was performed with immediate procedural success. At 6 month follow-up recurrent hyperhidrosis was reported affecting both axillae. A starch-iodine test showed a rim of persisting sweating at the periphery of both axillae.

Discussion: Curettage is generally performed without specific identification of sweat gland distribution, perhaps explaining the recurrence in this patient. In contrast, a starch-iodine test is used to guide administration of BOTOX[®] for hyperhidrosis. We propose pre-operative use of this technique before axillary curettage.

doi:10.1016/j.ejvs.2012.02.030

DOI of original article: 10.1016/j.ejvsextra.2012.02.008

Available online 12 May 2012

A Case of Iatrogenic Ilio-iliac Arteriovenous Fistula Initially Mistaken for Deep Venous Thrombosis

X. Liu^{a,b,c}, M. Lu^{a,b,c}, H. Shi^{a,b,c}, M. Jiang^{a,b,c}
^a *Department of Vascular Surgery, Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200011, China*
^b *Vascular Center, Shanghai Jiao Tong University School of Medicine, Shanghai 200011, China*
^c *Institute of Traumatic Medicine, Shanghai Jiao Tong University School of Medicine, Shanghai 200011, China*

Deep venous thrombosis (DVT) and associated complications cause significant morbidity and mortality in orthopedic surgery. Typical DVT symptoms, such as swelling, pain and discoloration in the affected extremities are often unreliable for diagnosis. Here we report a rare case of iatrogenic ilio-iliac arteriovenous fistula (AVF) due to lumbar discectomy, which was initially misdiagnosed as DVT, resulting in unnecessary implantation of a permanent inferior vena cava filter. Endovascular treatment is an attractive treatment option for such an AVF. We recommend a thorough physical and ultrasonography for patients presenting with DVT-like symptoms, especially following lumbar spinal surgery, to prevent overlooking underlying AVF.

doi:10.1016/j.ejvs.2012.03.019

DOI of original article: 10.1016/j.ejvsextra.2012.03.003

Available online 21 April 2012

[☆] Full articles available at www.ejvsextra.com